

Manifestation of a gap due to the exchange energy in a spinor condensate

Patrick Navez

*Katholieke Universiteit Leuven, Celestijnlaan 200 D, Heverlee, Belgium
Universitaet Duisburg-Essen, Lotharstrasse 1, 47057 Duisburg, Germany*

We investigate the dynamic response of population transfer between two components of a finite temperature spinor Bose condensed gas to a time-dependent coupling potential. Comparison between results obtained in the Bogoliubov-Popov approximation (BPA) and in the generalized random phase approximation (GRPA) shows noticeable discrepancies. In particular, the inter-component current response function calculated in the GRPA displays a gapped spectrum due to the exchange interaction energy whereas the corresponding density response function is gapless. We argue that the GRPA is better since, contrary to the BPA, it preserves the $SU(2)$ symmetry and the f -sum rule associated to the spinor gas. In order to validate the approximation, we propose an experimental setup that allows the observation of the predicted gap.

PACS numbers: 03.75.Hh, 03.75.Kk, 05.30.-d

I. INTRODUCTION

A great achievement in the discovery of condensation of Bose gas has been the experimental observation of the gapless and phonon-like nature of the collective excitations [1–3]. The experimental measurement of the energy spectrum as a function of momentum has confirmed the validity of the Bogoliubov theory [4]. However, well understood at zero temperature, the finite temperature extension of this theory is still controversial. The main difficulty is to find an approximation both *conserving* and *gapless* [5]. The generalized random phase approximation (GRPA) is such an approach that has been introduced first in the context of a quantum plasma in order to explain the collective phenomena [6]. Its application to Bose condensation has been successful to determine the sound velocity, the low-lying excitation spectrum observed in a trapped Bose gas [7] and the critical velocity [9]. Nevertheless, this theory suffers from the apparent contradiction that its building blocks contain propagators describing the particle motion in terms of the gapped and parabolic Hartree-Fock (HF) mean field dispersion relation [7, 8]. This paradox is resolved if the theory is reinterpreted adequately by making the distinction between collective excitations that result from oscillation of the condensate and quasi-particles or thermal excitations that compose the normal part of the gas [9]. Generally, in other fluids the two kinds of excitation exist (e.g. the plasmons and the electrons in a plasma). Therefore, no reason justifies that they should be identical in a Bose condensed gas as predicted from the Bogoliubov theory.

The only difference is that, in other fluids, the pole of the one particle Green function corresponds to the single particle spectrum while the pole of the susceptibility function corresponds to the collective excitations. In a Bose gas, below the condensation point, the two poles mix to form new poles that are identical for the two functions [5, 7]. In the non-conserving Bogoliubov theory, only one branch of excitation appears as a pole which corresponds to the quasi-particle resulting from hybridization of quantum states. In the GRPA, however,

the two branches of the susceptibility function and the branch of the one particle Green function mix to form three common branches of excitations, one of which corresponds to the phonon-like excitation [7]. In that case, it is not possible to associate a single quasi-particle to the one particle Green function with three branches of excitation as poles. For these reasons, the idea that the normal gas is composed of Bogoliubov-like quasi-particles seems too simplistic and must be reconsidered in the framework of a conserving approach. Basically, the GRPA tells that the excitation spectrum of a condensed particle is phonon-like while the one for the thermal particle is parabolic-like with a gap [8, 9]. The whole difficulty is how to perturb the Bose gas in order to selectively address these excitations.

In this paper, we aim at verifying the validity of these hypothetical thermal excitations by proposing an experiment on a spinor Bose gas revealing the existence of the gap. After reminding about the GRPA theory in section 2, we study in section 3 the inter-component transition between two internal levels of the Bose gas subjected to an external coupling field [10–13]. In particular, we are interested in determining the energy required to transfer a thermal atom from a Bose condensed gas towards another internal sub-level. According to the GRPA, such an atom becomes distinguishable from the condensed atoms and consequently releases a fixed amount of energy due to the Fock exchange interaction. An analog experiment is encountered in a superconductor-metal junction in which the measure of the electrical current as function of the voltage allows the determination of the BCS gap [14]. Both experimental [11] and theoretical studies [10] have been carried out on transitions between two internal levels in order to probe the inter-component density response function. But an external coupling has the effect to rotate the whole gas in the spinor space at no energy cost in the long wavelength limit and thus this function is gapless. For this reason, we investigate instead the inter-component current response function in order to be able to address separately the thermal atoms from the condensed ones and to display the gap. For com-

parison, a similar study is carried out in the gapless but non conserving Bogoliubov-Popov approximation (BPA) [4, 5]. This approach extends the Bogoliubov theory to strongly depleted gas and leads to different results. In section 4, we generalize the approach to the case of real experiments with a trap while section 5 ends with the conclusion.

II. THE GRPA FORMALISM

To start with, we consider a spinor Bose condensed gas of atoms with mass m in a volume Ω populated only in two sub-levels chosen among the hyperfine structure of the atom and labeled by $\alpha = 1, 2$. The scattering lengths between atoms of any sub-level are assumed to be equal to a (in ^{87}Rb their relative difference is about 10^{-2} [12]) and define the coupling constant $g = 4\pi a/m$ ($\hbar = 1$). Defining the annihilation operators $c_{\alpha,\mathbf{k}}$ where $\mathbf{k}$ is the momentum and $\epsilon_{\mathbf{k}} = \mathbf{k}^2/2m$, the Hamiltonian is written as:

$$H = \sum_{\alpha,\mathbf{k}} \epsilon_{\mathbf{k}} c_{\alpha,\mathbf{k}}^\dagger c_{\alpha,\mathbf{k}} + \frac{g}{2\Omega} \sum_{\alpha,\beta,\mathbf{k},\mathbf{k}',\mathbf{q}} c_{\alpha,\mathbf{k}}^\dagger c_{\beta,\mathbf{k}'}^\dagger c_{\alpha,\mathbf{k}+\mathbf{q}} c_{\beta,\mathbf{k}'-\mathbf{q}} \quad (1)$$

The eventual energy shift between the two sub-levels is removed by choosing an appropriate representation of $c_{\alpha,\mathbf{k}}$. We shall study the dynamic evolution of the excitation operator $\rho_{\mathbf{k},\mathbf{q}}^{\alpha,\beta}(t) = e^{iHt} c_{\alpha,\mathbf{k}}^\dagger c_{\beta,\mathbf{k}+\mathbf{q}} e^{-iHt}/\Omega$ in terms of which we express the space Fourier transform of the densities and the currents, respectively $\rho_{\mathbf{q}}^{\alpha,\beta}(t) = \sum_{\mathbf{k}} \rho_{\mathbf{k},\mathbf{q}}^{\alpha,\beta}(t)$ and $\mathbf{J}_{\mathbf{q}}^{\alpha,\beta}(t) = \sum_{\mathbf{k}} \rho_{\mathbf{k},\mathbf{q}}^{\alpha,\beta}(t)(\mathbf{k} + \mathbf{q}/2)/m$. These are associated to the $U(1)$ number conservation and $SU(2)$ spinor symmetries of H . We first consider the case of a uniform condensed gas initially stable populated in the sub-level 1 such that $\langle \rho_{\mathbf{k},\mathbf{q}}^{\alpha,\beta} \rangle = \delta_{\alpha,1} \delta_{\alpha,\beta} \delta_{\mathbf{q},0} n_{1,\mathbf{k}}$. $\langle \dots \rangle$ denotes the statistical ensemble average $\text{Tr}(\sigma \dots)$ where σ is the density matrix, $n_{1,\mathbf{k}}$ the momentum density in the mode $\mathbf{k}$ and $n_c = n_{1,0}$ the condensate density.

Let us introduce a coupling between the two spinor components through the time dependent Hamiltonian $H_V(t) = \Omega \rho_{\mathbf{q}}^{12\dagger} V_{\mathbf{q},\omega}^{12} e^{-i(\omega+i0)t} + c.c.$ where $V_{\mathbf{q},\omega}^{12}$ is a constant and $i0$ ensures the adiabatic condition. The change in the frequency mode ω of the inter-component density $\delta \rho_{\mathbf{q},\omega}^{12}$ from its equilibrium value $\langle \rho_{\mathbf{q}}^{1,2} \rangle = 0$ induced by this potential is related to the inter-component susceptibility function through:

$$\delta \rho_{\mathbf{q},\omega}^{12} = \chi_{12}(\mathbf{q}, \omega) V_{\mathbf{q},\omega}^{12} \quad (2)$$

Formal linear response theory allows to find the formula [15]:

$$\chi_{12}(\mathbf{q}, \omega) = i\Omega \int_0^\infty dt e^{i(\omega+i0)t} \langle [\rho_{\mathbf{q}}^{12\dagger}(0), \rho_{\mathbf{q}}^{12}(t)] \rangle \quad (3)$$

Although the initial state is not the $SU(2)$ symmetric equilibrium state, the linear response formalism is valid

in a much more general context of a stable state such that $[H, \sigma] = 0$. Similarly to the intra-component case, provided that this function is causal and the $n_{\alpha,\mathbf{k}}$'s are isotropic, its imaginary part obeys the following f -sum rule [10]:

$$\int_{-\infty}^\infty d\omega \omega \text{Im} \chi_{12}(\mathbf{q}, \omega) = -\pi \frac{\mathbf{q}^2}{2m} n_1 \quad (4)$$

This function is also connected to the longitudinal current response function $\chi_{12}^{\parallel}(\mathbf{q}, \omega)$ through the exact relation:

$$\chi_{12}(\mathbf{q}, \omega) = \frac{n_1}{\omega + i0} + \frac{\mathbf{q}^2}{2m} \frac{n_1}{(\omega + i0)^2} + \frac{\mathbf{q}^2 \chi_{12}^{\parallel}(\mathbf{q}, \omega)}{(\omega + i0)^2} \quad (5)$$

This last function is defined by replacing $\rho_{\mathbf{q}}^{12}(t)$ by $\mathbf{q} \cdot \mathbf{J}_{\mathbf{q}}^{12}(t)/|\mathbf{q}|$ in (3). The f -sum rule and Eq.(5) result from the conservation law $\partial_t \rho_{\mathbf{q}}^{12}(t) = i\mathbf{q} \cdot \mathbf{J}_{\mathbf{q}}^{12}(t)$ associated to the $SU(2)$ symmetry.

The GRPA approach is based on the assumption that the off-diagonal part of the excitation operator ($\mathbf{q} \neq 0$) contributes less than the diagonal one ($\mathbf{q} = 0$) because the operators involved do not oscillate in phase but rather randomly [9]. As a consequence, in the Heisenberg equation for $\rho_{\mathbf{k},\mathbf{q}}^{12}(t)$, we neglect all contributions quadratic in the off-diagonal operator in comparison with products involving off and diagonal ones. In this way, we obtain an equation of motion of a similar form to the intra-component case [9, 10]:

$$(i\partial_t - \epsilon_{2,\mathbf{k}+\mathbf{q}}^{HF} + \epsilon_{1,\mathbf{k}}^{HF}) \rho_{\mathbf{k},\mathbf{q}}^{12}(t) = g n_{1,\mathbf{k}} \rho_{\mathbf{q}}^{12}(t) \quad (6)$$

where $\epsilon_{\alpha,\mathbf{k}}^{HF} = \epsilon_{\mathbf{k}} + g n_1 (1 + \delta_{1,\alpha})$ is the HF energy felt by the atom in sub-level α .

III. RESULTS FOR THE HOMOGENEOUS GAS

The operatorial integral equation (6) is solved exactly using methods developed in [9]. Plugging the solution into (3), the susceptibility function is found to be [8, 10]:

$$\chi_{12}(\mathbf{q}, \omega) = \chi_{0,12}(\mathbf{q}, \omega) / (1 - g \chi_{0,12}(\mathbf{q}, \omega)) \quad (7)$$

where

$$\chi_{0,12}(\mathbf{q}, \omega) = \sum_{\mathbf{k}} \frac{n_{1,\mathbf{k}}}{\omega + i0 + \epsilon_{1,\mathbf{k}}^{HF} - \epsilon_{2,\mathbf{k}+\mathbf{q}}^{HF}} \quad (8)$$

As shown in [10], this function is gapless since the pole $\omega = 0$ corresponds to a rotation in the $SU(2)$ space. Using the Kramers-Kronig relations [4], we check that Eq.(7) fulfills the f -sum rule (4). Also, by introducing a vector potential as an external perturbation, the longitudinal current response function can be calculated as well in this approximation and the result verifies the property (5). Combination of (5), (7) and (8) in the limit of long wavelength allows to obtain:

$$\chi_{12}^{\parallel}(\mathbf{q} = 0, \omega) = \frac{\sum_{\mathbf{k}} 2\epsilon_{\mathbf{k}} n_{1,\mathbf{k}} / (3m)}{\omega + i0 + g n_1} \quad (9)$$

We find a non zero pole $-gn_1$ that corresponds to a gap due to the Fock exchange interaction and that is interpreted as the transition potential energy for any transferred thermal atom. The fluctuations of the inter-component longitudinal current induce a relative motion between the normal fluid and the superfluid in the spin space. Note that, contrary to the intra-component case, the longitudinal current response function Eq.(9) does not have the same pole as the density response function Eq.(7) and is identical to the transverse current response function when $\mathbf{q} \rightarrow \mathbf{0}$. Eq.(9) is the main result of this paper and we shall show how this gap can be observed in experiments with a trap.

In contrast, the use of the BPA leads to drastically different results. It corresponds to the Hartree-Fock-Bogoliubov approximation to the Hamiltonian in which the term quadratic in the product of anomalous average is also neglected [4]. The chemical potential is given by $\mu = g(2n_1 - n_c)$, the quantum operators describing the various components evolve according to $c_{1,\mathbf{k}}(t) = e^{-i\mu t}(\sqrt{\Omega n_c} \delta_{\mathbf{k},0} + u_{+,\mathbf{k}} e^{-i\epsilon_{\mathbf{k}}^B t} b_{1,\mathbf{k}} + u_{-,\mathbf{k}} e^{i\epsilon_{\mathbf{k}}^B t} b_{1,-\mathbf{k}}^\dagger)$ and $c_{2,\mathbf{k}}(t) = e^{-i(\mu + \epsilon_{\mathbf{k}})t} c_{2,\mathbf{k}}$. $b_{1,\mathbf{k}}$ is the annihilation operator associated to the quasi-particle, $\epsilon_{\mathbf{k}}^B = \sqrt{2gn_c \epsilon_{\mathbf{k}} + \epsilon_{\mathbf{k}}^2}$ and $u_{\pm,\mathbf{k}} = \pm((\epsilon_{\mathbf{k}} + gn_c)/2\epsilon_{\mathbf{k}}^B \pm 1/2)^{1/2}$. As a result, using (3), we obtain:

$$\chi_{12}^{BP}(\mathbf{q}, \omega) = \frac{n_c}{\omega - \epsilon_{\mathbf{q}} + i0} + \sum_{\pm, \mathbf{k}} \frac{u_{\pm, \mathbf{k}}^2 (n_{1, \mathbf{k}}^B + \delta_{\pm, -})}{\omega + i0 \pm \epsilon_{\mathbf{k}}^B - \epsilon_{\mathbf{k} \pm \mathbf{q}}} \quad (10)$$

$$\chi_{12}^{\parallel BP}(\mathbf{0}, \omega) = \sum_{\pm, \mathbf{k}} \frac{2\epsilon_{\mathbf{k}}}{3m} \frac{u_{\pm, \mathbf{k}}^2 (n_{1, \mathbf{k}}^B + \delta_{\pm, -})}{\omega + i0 \pm \epsilon_{\mathbf{k}}^B - \epsilon_{\mathbf{k}}} \quad (11)$$

where $n_{1, \mathbf{k}}^B = \langle b_{1, \mathbf{k}}^\dagger b_{1, \mathbf{k}} \rangle$. Both approaches GRPA and BPA are gapless and their susceptibilities defined respectively by (7) and (10) have a pole at $\omega = \epsilon_{\mathbf{q}}$ for zero temperature in agreement with the spectrum obtained from the linearization of the coupled Gross-Pitaevskii equations [16]. Besides this pole, we note from Eq.(10-11) that transitions in BPA result from scattering of a phonon into a normal atom releasing energy $\epsilon_{\mathbf{k}}^B - \epsilon_{\mathbf{k} + \mathbf{q}}$ and from creations of both a phonon and a normal atom absorbing the energy $\epsilon_{\mathbf{k}}^B + \epsilon_{\mathbf{k} - \mathbf{q}}$. As a consequence, for $\mathbf{q} = 0$, the transition frequency spectrum is continuous with a line-width estimated to gn_1 . In GRPA, however, only a process that releases the constant gap energy happens without any line-width.

Therefore, the two theories display noticeable differences but strong theoretical arguments play in favor of GRPA. Due to the $SU(2)$ breaking of symmetry, the BPA violates the conservation laws and thus does not obey the f -sum rule nor Eq.(5). In particular, for $\mathbf{q} = 0$ and $\omega = 0$, the action of the external coupling is to rotate the whole gas in the spin space while it rotates only the condensate in the BPA. Physically, this violation can be understood by noticing that a phonon has not a well defined atom number ($[b_{1, \mathbf{k}}, \rho_{\mathbf{k}, 0}^{11}] \neq b_{1, \mathbf{k}}$) and therefore cannot be transformed into an atom alone during a transition.

For these reasons, the GRPA is a better approximation that is justified if the predicted gap is observed in experiments.

IV. RESULTS FOR THE TRAPPED GAS

The considerations so far obtained are straightforwardly extended to a trapped gas. We start with a condensed gas initially populated in the sub-level 1 confined in a parabolic potential $V^{11}(\mathbf{r}) = \sum_{i=x,y,z} \frac{1}{2} m \omega_i^2 r_i^2$. In this new situation, the density profile $n_1(\mathbf{r})$ depends on the position and is determined in good approximation from the mean field HF equations in the Thomas-Fermi limit [4]: the condensate density $n_c(\mathbf{r})$ is determined from the Gross-Pitaevskii equation in absence of the kinetic term $\mu = V^{11}(\mathbf{r}) + g(2n_1(\mathbf{r}) - n_c(\mathbf{r}))$; the thermal atoms density is determined from the semi-classical expression: $n_{1, \mathbf{k}}(\mathbf{r}) = 1/[\exp(\beta(\epsilon_{\mathbf{k}}^{HF}(\mathbf{r}) + V^{11}(\mathbf{r}) - \mu)) - 1]$ where $\beta = 1/k_B T$ is the inverse temperature. Would we have applied the BPA, we should have rather used $n_{1, \mathbf{k}}^B(\mathbf{r}) = 1/(\exp(\beta \epsilon_{\mathbf{k}}^B(\mathbf{r})) - 1)$ in the condensate region. We assume an isotropic trap with frequency $\omega_i = \bar{\omega} = 2\pi \times 100 \text{ Hz}$, scattering length $a\sqrt{m\bar{\omega}} = 10^{-3}$ and $\mu = 2\pi \times 2k \text{ Hz}$. In the second sub-level, we consider both cases of a trapped and a flat potential $V^{22}(\mathbf{r}) = V^{11}(\mathbf{r}), 0$. In order to induce a current in the trap and transfer atoms to the second sub-level, we apply the external coupling $V^{12}(\mathbf{r}, t) = \mathbf{r}_z \cdot \mathbf{F}_z e^{i\omega_0 t}$ during a finite time interval and beginning at $t = 0$ where $\mathbf{F}_z$ refers to a force constant along the z axis in the real space. This coupling can be produced either by a rotating magnetic field gradient $\mathbf{B} = B'r_z(\cos(\omega_0 t), \sin(\omega_0 t), 0)$ that interacts with the gas with its spin orientation along the z -axis [12], or by a two photon Raman scattering between two multiplets (e.g. $F = 1$ and $F = 2$ in ^{87}Rb) creating a stationary optical field with a node centered at the minimum of the trap [13].

In presence of inhomogeneity, the density response has to be generalized into [3, 4]:

$$\delta\rho^{12}(\mathbf{r}_1, t) = \int_0^t dt' \int_\Omega d^3\mathbf{r}_2 \chi_{12}(\mathbf{r}_1, \mathbf{r}_2, t - t') V^{12}(\mathbf{r}_2, t') \quad (12)$$

The total number of output atoms in the second level $N_2 = \int_\Omega d^3\mathbf{r} n_2(\mathbf{r}, t)$ is determined through the transfer rate equation:

$$\frac{dN_2}{dt} = i \int_\Omega d^3\mathbf{r}_1 V^{21}(\mathbf{r}_1, t) \delta\rho^{12}(\mathbf{r}_1, t) + c.c. \quad (13)$$

For $t \gg 1/\omega_0$, the time integral can be extended to infinity in (12) and Eq.(13) becomes time-independent. Combining Eqs.(12-13), rewriting $\mathbf{r}_1 = \mathbf{r} + \mathbf{r}'/2$ and $\mathbf{r}_2 = \mathbf{r} - \mathbf{r}'/2$ and using $\chi_{12}(\mathbf{r}, \mathbf{q}, \omega) = \int_0^\infty dt \int_\Omega d^3\mathbf{r}' e^{-i(\mathbf{q} \cdot \mathbf{r}' + \omega t)} \chi_{12}(\mathbf{r}_1, \mathbf{r}_2, t)$, the transfer rate is expressed up to the second order in the coupling in terms of the inhomogeneous susceptibility $\chi_{12}(\mathbf{r}, \mathbf{q}, \omega)$.




FIG. 1: In (a), schematic representation of the local potential energy (full line) in a trap felt by thermal atoms for the trapped sub-level 1 (left) and 2 (right). The difference accounts for the transition energy. In (b), corresponding rate of output atoms in the second sub-level in GRPA (black) and in BPA (grey) for $T/T_c = 0.56$.

This last function is calculated in the local density approximation by replacing the density and the frequency in the homogeneous expressions (5) and (9) by their local forms $n_1(\mathbf{r})$ and $\omega(\mathbf{r}) = V^{11}(\mathbf{r}) - V^{22}(\mathbf{r}) - \omega_0$ [3]. This procedure gives successively:

$$\begin{aligned} \frac{1}{F_z^2} \frac{dN_2}{dt} &= -2\pi \int d^3\mathbf{r} \left(r_z^2 + \frac{\partial_{q_z}^2}{4} \right) \text{Im} \chi_{12}(\mathbf{r}, \mathbf{q}, \omega_0)|_{\mathbf{q} \rightarrow 0} \\ &= -2\pi \int d^3\mathbf{r} \text{Im} \left[\frac{r_z^2 n_1(\mathbf{r})}{\omega(\mathbf{r}) + i0} + \frac{1}{(\omega(\mathbf{r}) + i0)^2} \right. \\ &\quad \times \left. \left(\frac{n_1(\mathbf{r})}{4m} + \frac{\sum_{\mathbf{k}} \epsilon_{\mathbf{k}} n_{1,\mathbf{k}}(\mathbf{r})/(3m)}{\omega(\mathbf{r}) + i0 + gn_1(\mathbf{r})} \right) \right] \end{aligned} \quad (14)$$

Eq.(14) is expected to describe accurately the transfer rate provided $\omega_0 \gg \bar{\omega}$ and $t \ll \bar{\omega}^{-1}$ so that the effects of discretization in the excitation spectrum can be safely ignored. These effects also diminish due to the higher density states at finite temperature. Also, the linear response theory is valid provided that the typical time duration t is much lower than the inverse of the typical Rabi frequency estimated to $F_z R_{TF}$ where R_{TF} is the condensate size. This implies the upper bound $F_z \ll \bar{\omega}/R_{TF} \sim \bar{\omega}^2 \sqrt{m/2\mu}$. Thus, in GRPA, the total number of transferred atom results from a local rotation of the gas in the spin space in regions where $\omega_0 = V^{11}(\mathbf{r}) - V^{22}(\mathbf{r})$ and from an inter-component current of thermal atoms releasing the exchange Fock interaction energy in regions where $\omega_0 = gn_1(\mathbf{r}) + V^{11}(\mathbf{r}) - V^{22}(\mathbf{r})$.

In the case of $V^{22}(\mathbf{r}) = V^{11}(\mathbf{r})$, rotation in the spin space occurs only for zero frequency so that a non zero frequency rate provides a direct evidence of the gap $\omega_0 = gn_1(\mathbf{r})$. This released energy depends mainly on the local condensate density (see Fig.1a). Thus the transfer




FIG. 2: In (a), same as Fig.1 but for a transition to an untrapped sublevel. Note the constant gap in the condensate region responsible for the peak. Corresponding rate of output atoms for $T/T_c = 0.56$ in (b) and $T/T_c = 0.85$ in (c).

$R = m(\bar{\omega}/F_z)^2 dN_2/dt$ calculated numerically from (14) and shown in Fig.1b as a function of ω_0 displays an inhomogeneous broadening. This rate decreases with frequency whereas the density of thermal atoms is higher in the edge of the condensate than in the center of the trap. For comparison, we plot the transition rate obtained in BPA by inserting Eq.(10) in Eq.(14) where contrary to GRPA energy absorption is possible when $\omega_0 < 0$. The reduced transition rate results from the broad spectrum displayed in Eq.(10).

Fig.2 shows the same curves as Fig.1 but for the case $V^{22}(\mathbf{r}) = 0$. The output atoms are not trapped in order to diminish the dispersion due to the inhomogeneity. As a result, a resonant transition occurs for a frequency close to $\omega_0 = gn_1(\mathbf{r} = 0)$. However the transition due to the spinor rotation becomes frequency dependent and leads to a background noise. The broadening of the resonance is caused by the variation of the thermal density in the condensate region. The line-width is estimated to $\Delta\omega \sim g(mk_B T/2\pi)^{3/2}$ and increases with the temperature. Such a broader peak is observed in Fig.2c but at the expense of a more intense background noise. This noise

is proportional to the local gas inertia $r_z^2 n_1(\mathbf{r})$ about the z-axis (see Eq.(14)) and is more important for stronger depletion in the region of high frequency. It considerably reduces if we choose $\omega_z \gg \bar{\omega}$ without altering the validity of the local density approximation ($\omega_z \ll \omega_0$).

V. CONCLUSIONS

In conclusion, we analyze theoretically the atom transition rate between two sub-levels of a spinor condensate

caused by a time dependent external coupling. Significant differences are reported between the GRPA and BPA models. An experimental setup is suggested in order to validate at most one approach and can serve to understand the true nature of the quasi-particle dispersion relation.

PN thanks K. Bongs for usefull discussions and acknowledges support from the Belgian FWO project G.0115.06, from the Junior fellowship F/05/011 of the KUL research council, and from the German AvH foundation.

-
- [1] D.M. Stamper-Kurn, A.P. Chikkatur, A. Görlitz, S. Inouye, S. Gupta, D.E. Pritchard, and W. Ketterle, Phys. Rev. Lett. **83**, 2876 (1999).
 - [2] N.M. Hugenholtz and D. Pines, Phys. Rev. **116**, 489 (1959).
 - [3] F. Zambelli, L. Pitaevskii, D.M. Stamper-Kurn and S. Stringari, Phys. Rev. A **61**, 063608 (2000); R. Ozeri, N. Katz, J. Steinhauer, and N. Davidson, Rev. Mod. Phys. **77**, 187 (2005).
 - [4] A. J. Leggett, Rev. Mod. Phys. **73**, 307 (2001); L. Pitaevskii and S. Stringari, *Bose-Einstein Condensation*, (Clarendon Press, 2003); C. J. Pethick and H. Smith, *Bose-Einstein Condensation in Dilute Gases*, (Cambridge University Press, 2001).
 - [5] A. Griffin, Phys. Rev. B **53**, 9341 (1996); P.C. Hohenberg and P.C. Martin, Ann. Phys. (N.Y.) **34**, 291 (1965); A. Griffin, Excitations in a Bose-Condensed Liquid (University Press, Cambridge, 1993).
 - [6] D. Pines and P. Nozières, *The theory of Quantum Liquids*, vol 1 (Benjamin, New York, 1966).
 - [7] P. Szépfalusy and I. Kondor, Ann. Phys. **82**, 1-53 (1974); M. Fliesser, J. Reidl, P. Szépfalusy, R. Graham, Phys. Rev. A **64**, 013609 (2001); J. Reidl, A. Csordas, R. Graham, P. Szépfalusy, Phys. Rev. A **61**, 043606 (2000).
 - [8] C-H Zhang and H.A. Fertig, Phys. Rev. A **75**, 013601 (2007).
 - [9] P. Navez, J. Low. Temp. Phys. **138**, 705-710 (2005); P. Navez, Physica A **356**, 241-278 (2005); P. Navez and R. Graham, Phys. Rev. A **73**, 043612 (2006).
 - [10] M.Ö. Oktel and L.S. Levitov, Phys. Rev. Lett. **83**, 1 (1999); M.Ö. Oktel and L.S. Levitov, Phys. Rev. A **65**, 063604 (2002); M.Ö. Oktel, T.C. Killian, D. Kleppner and L.S. Levitov, Phys. Rev. A **65**, 033617 (2002); M.Ö. Oktel and L.S. Levitov, Phys. Rev. Lett. **88**, 230403 (2002).
 - [11] D.G. Fried, T.C. Killian, L. Willmann, D. Landhuis, S.C. Moss, D. Kleppner and T.J. Greytak, Phys. Rev. Lett. **81**, 3811 (1998).
 - [12] J. M. Higbie, L. E. Sadler, S. Inouye, A. P. Chikkatur, S. R. Leslie, K. L. Moore, V. Savalli, and D. M. Stamper-Kurn, Phys. Rev. Lett. **95**, 050401 (2005); J. Kronjäger, C. Becker, M. Brinkmann, R. Walser, P. Navez, K. Bongs, and K. Sengstock Phys. Rev. A **72**, 063619 (2005).
 - [13] H.J. Lewandowski, J. M. McGuirk, D. M. Harber and E. A. Cornell, Phys. Rev. Lett. **91**, 240404 (2003).
 - [14] For example see R.P. Feynmann, *Statistical Mechanics* (Perseus Books groups, 1972).
 - [15] R. Balescu, *Equilibrium and non equilibrium statistical mechanics* (John Wiley & Sons, New York, 1975).
 - [16] W. B. Colson and A. L. Fetter, J. Low Temp. Phys. **33**, 231 (1978); W. Zhang, D. L. Zhou, M.-S. Chang, M. S. Chapman, and L. You, Phys. Rev. Lett. **95**, 180403 (2005).